



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
NEW JERSEY

ABOUND[®] FLOWABLE FUNGICIDE

EPA Reg. No. 100-1098
EPA SLN No. NJ-050002

To allow fungicide applications to begin prior to bloom and for suppression of
fairy ring disease (*Psilocybe sp.*) in cranberries

Active Ingredient:

Azoxystrobin: methyl (E)-2-{2-[6-(2-cyanophenoxy)

Pyrimidin-4-yloxy]phenyl]-3-methoxyacrylate -----22.9%

Other Ingredients: -----77.1%

Total: -----100.0%

Contains 2.08 lbs. Of active ingredient per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS
LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY OR ILLEGAL
RESIDUES.

It is a violation of Federal law to use this product in a manner inconsistent with its
labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements
and precautions, on the EPA-registered Abound Flowable Fungicide label.

This label must be in the possession of the user at the time of pesticide mixing and
application.

DIRECTIONS FOR USE

| | | | |
|---|--|--------------------------------|--|
| Cranberry | <p>Cottonball (<i>Monilia oxycocci</i>)</p> <p>Lophodermium twig blight</p> <p>Fruit rots (<i>Glomerella cingulata</i>) (<i>Coleophoma empetri</i>)</p> | <p>6.2-15.4 (0.1-0.25)</p> | <p>Integrated Pest (Disease) Management: ABOUND should be integrated into an overall disease management strategy that includes selection of varieties with disease tolerance, optimum plant populations, proper fertilization, plant residue management, crop rotation and proper water management.</p> <p>Resistance Management: Do not apply more than two sequential applications of ABOUND or other QoI fungicides before alternating with a fungicide that has a different mode of action. Do not make more than six applications of ABOUND or other QoI fungicides per crop per acre per year.</p> <p>Application Directions: Begin applications at the onset of disease but prior to 5-10% bloom for fruit rot, cottonball, and twig blight. Continue applications on a 7-14 day schedule if conditions are favorable for disease development. Applications may be made by ground, chemigation or air.</p> |
| | <p>Fairy ring (suppression) (<i>Psilocybe</i> sp)</p> | <p>15.4 (0.25)</p> | <p>Application Directions: Make the first application at bud break. Measure the ring diameter and add 10 feet to diameter. Apply ABOUND at a rate equivalent to 15.4 oz/acre in 30-100 gallons of water to the affected area. Irrigation (1-2 hours) following application is advisable to ensure penetration to the base of the plant. Repeat 2-4 weeks later, if necessary. For ground application ensure adequate water volume for thorough canopy penetration.</p> |
| <p>Specific Use Restrictions: May be applied up to three days prior to harvest (3-day PHI). Do not apply more than 2.88 quarts product (1.5 lb ai) per acre per season. Do not treat cranberry fields used for aquaculture of fish and crustacea Do not apply when weather conditions favor drift from treated areas to non-target aquatic habitat. Applicators should use care in applications near non-target aquatic habitats. Do not apply to flooded crop. Do not allow release of irrigation or flood water to non-target aquatic habitat for at least 14 days after the last application.</p> | | | |

Abound is registered trademark of Syngenta

24c registrant:
 Syngenta Crop Protection, Inc
 PO Box 18300
 Greensboro, NC 27419-8300

5/22/12 for n - name of Syngenta

1/1

Label Code: NJ1098xxxAA0305

Abound Flowable Fungicide/Cranberry/NJ-050002