



Bayer CropScience

Bayer CropScience LP  
 P.O. Box 12014  
 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

**Admire® 2 Flowable Insecticide**

**For Use on Christmas Trees.**

EPA Reg. No. 264-758  
 EPA SLN. No. NJ-050001

## ***24(c) Supplemental Label***

### **FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW JERSEY**

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label and the EPA label must be in the possession of the user at the time of pesticide mixing and application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Do not apply within 25 feet of lakes, reservoirs, permanent streams, marshes, natural ponds, estuaries and commercial farm ponds. Do not cultivate within 10 feet of any of these areas to allow growth of a vegetative filter strip. When using this product on any erodible soils, Best Management Practices for minimizing runoff should be employed. Consult your local Natural Resources Conservation Service for recommendations in your use area.

#### **CHRISTMAS TREE**

<b>Pests Controlled</b>	<b>Rate fluid ounces/Acre</b>
White grub complex (damage from grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and oriental beetle)	16.0 – 32.0
<b>Notes and Restrictions</b> Maximum ADMIRE allowed per season: 32.0 fluid ounces/Acre (0.5 lb AI/Acre). <b>Applications</b> Soil incorporation and movement of ADMIRE to the root zone is required for activity. ADMIRE can be incorporated most readily when applied to moist soil. Apply specified dosage of ADMIRE in one of the following methods: 1. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. 18-inch band on each side of the row (small trees) to full broadcast application (large trees) followed by rainfall or 0.25 - 1 inch of irrigation within 24 hours after application.  For optimal grub control, apply ADMIRE during adult flight activity, or up to mid-July, when 1st instar larvae are present.	

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

*As with any crop-protection product, always read and follow label directions.  
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*